

Drug Name: Nucala (mepolizumab)

Revised Date: 12/2018

Drug Name:	Nucala (mepolizumab)
Exclusion	n/a
Criteria:	
Required Medical Information:	
	<ul> <li>Eosinophilic Asthma</li> <li>Authorization of 12 months may be granted for treatment of eosinophilic asthma when all of the following criteria are met: <ul> <li>Member is 12 years of age or older.</li> <li>Member has a baseline blood eosinophil count of at least 150 cells per microliter.</li> <li>Member has a history of severe asthma despite current treatment with both of the following medications at optimized doses: <ul> <li>Inhaled corticosteroid</li> <li>Additional controller (long acting beta<sub>2</sub>-agonist, leukotriene modifier, or sustained-release theophylline)</li> </ul> </li> </ul></li></ul>
	<ul> <li>Eosinophilic Granulomatosis with Polyangiitis</li> <li>Authorization of 12 months may be granted for treatment of eosinophilic granulomatosis with polyangiitis when all of the following criteria are met:</li> <li>Member is 18 years of age or older.</li> <li>Member has a history or the presence of an eosinophil count of more than 1000 cells per microliter or a blood eosinophil level of greater than 10%.</li> </ul>
Renewal Criteria	<ul> <li>Eosinophilic asthma</li> <li>Authorization of 12 months may be granted for continuation of treatment of eosinophilic asthma when all of the following criteria are met: <ul> <li>Member is 12 years of age or older.</li> <li>Asthma control has improved on Nucala treatment as demonstrated by either: <ul> <li>A reduction in the frequency or severity of symptoms and exacerbations, OR a reduction in the daily maintenance oral corticosteroid dose</li> </ul> </li> </ul></li></ul>
	Eosinophilic Granulomatosis with Polyangiitis  Authorization of 12 months may be granted for continuation of treatment of eosinophilic granulomatosis with polyangiitis when all of the following criteria are met:  • Member is 18 years of age or older.  • Member has beneficial response to treatment with Nucala as



## demonstrated by any of the following:

- 0 A reduction in the frequency of relapses, or
- o A reduction in the daily oral corticosteroid dose, or
- o No active vasculitis